


ICPIG 2017

XXXIII INTERNATIONAL CONFERENCE
ON PHENOMENA IN IONIZED GASES

CONFERENCE PROCEEDINGS



Proceedings of the
XXXIII
INTERNATIONAL
CONFERENCE ON
PHENOMENA IN IONIZED
GASES

Estoril, Portugal. 9-14 July 2017

Editors:

Luís Lemos Alves
Antonio Tejero-del-Caz

Published by:

Instituto de Plasmas e Fusão Nuclear

Instituto Superior Técnico

Universidade de Lisboa

<http://icpig2017.tecnico.ulisboa.pt>

Credits:

Editors: Luís Lemos Alves

Antonio Tejero-del-Caz

Cover design: Irene Lemos Alves

The XXXIII ICPIG (International Conference on Phenomena in Ionized Gases) has been organized by Instituto de Plasmas e Fusão Nuclear from Instituto Superior Técnico, Universidade de Lisboa, Universidade do Porto and Universidade do Minho.

The XXXIII ICPIG was organized in accordance with IUPAP principles, regarding the free circulation of scientists for international collaborations and discussions, as stated in the declaration of the International Council of Science, adopted at the 26th General Assembly in 2008 and endorsed by the 27th General Assembly in 2011. In particular, no *bona fide* scientist is excluded from participation on grounds of origin, nationality or political considerations unrelated to science.

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. Abstracting is permitted with credit to the source.

Bio-relevant NO_x generated by transient spark in atmospheric dry air and air with water electropray

Z. Machala¹, K. Hensel¹, B. Tarabová¹, M. Janda¹

¹ Faculty of Mathematics, Physics and Informatics, Comenius University in Bratislava, Slovakia

Generation of nitrogen oxides (NO_x) was studied in a DC-driven self-pulsing transient spark (TS) discharge in atmospheric pressure air. The precursors of NO_x production and the TS characteristics were studied by nanosecond time-resolved optical diagnostics. Thanks to the short (~20–50 ns) high current (~1 A) spark pulses, highly reactive non-equilibrium plasma is generated. The NO_x production rate of $\sim 7 \times 10^{16}$ molecules/J was achieved in dry air, dependent on TS repetition frequency, i.e. power, which is related to the complex frequency-dependent discharge properties and thus NO₂/NO generating mechanisms. With water electrosprayed through the TS, gaseous NO_x formation was lowered but induced chemical changes in water make it of biomedical importance.

1. Introduction

Nitrogen oxides (NO_x) are typical by-products of air plasmas that have important bio-relevant properties, e.g. as antimicrobial (NO₂), physiological (NO), and anesthetics (N₂O) agents. We studied the generation of NO and NO₂ in the transient spark (TS) discharge in atmospheric pressure air, using optical emission spectroscopy combined with the post-discharge gas composition analysis by FTIR.

The TS is a DC-driven repetitive self-pulsing discharge with 20-50 ns short spark current pulses initiated by streamers, with the pulse repetition frequency 1-10 kHz [1]. It has been successfully applied for flue gas cleaning and biodecontamination of water [2]. The air TS can be combined with the electropray of water, which induces formation of nitrites, nitrates, hydrogen peroxide and peroxyxynitrites and demonstrates strong antibacterial properties of such plasma activated water [2].

2. Results

Generation of NO_x in the gas phase was studied in dry air, and in the air humidified by water electrosprayed through the discharge. The dominant stable gas phase products in dry air were nitrogen oxides, while ozone was not detected (<10 ppm detection limit). NO formation steeply increases with the discharge power, as shown in Fig. 1. The sum of NO and NO₂ concentration >400 ppm was achieved with power input below 6 W. The highest NO_x (NO + NO₂) generation rate achieved was around 7×10^{16} molecules/J [3]. Due to their easy dissolution in the water and possibly also due to the discharge cooling by water and thus decreased NO_x formation, the NO_x densities were found lower in air humidified by the water electropray (Fig. 1).




Fig. 1: NO and NO₂ generation in TS in dry air vs. air with water electropray for increasing discharge power.

3. Summary

TS in atmospheric air provided high production rates of NO_x. With sprayed water, gaseous NO_x formation was lowered but induced chemical changes in water make it of biomedical importance.

3. References

- [1] M. Janda, V. Martišoviš, L. Dvonč, et al., *Plasma Sources Sci. Technol.*, **23**, 065016 (2014).
- [2] Z. Machala, B. Tarabová, K. Hensel, et al., *Plasma Process. Polym.*, **10**, 649 (2013).
- [3] M. Janda, V. Martišoviš, K. Hensel, Z. Machala, *Plasma Chem. Plasma Process.* **36**, 767 (2016)

This work was supported by Slovak Research and Development Agency APVV-0134-12 and Slovak grant agency VEGA 1/0918/15.